



## Post-Operative Instructions for LIP REPOSITIONING PROCEDURE

With proper attention to post-operative care, most dental surgeries would heal quickly and without complication;

- **Control of Bleeding:** A slight amount of blood mixed with your saliva is normal following any dental surgery and it may persist until the next day. Avoid rinsing or spitting during the first night. **Try to keep your head elevated while sleeping for the first couple of nights.** Later, if bleeding starts again, please elevate your head and apply the cold moist gauze or tea bag with gentle pressure over the surgical site for at least 20 minutes, and repeat if necessary. Should it persist severely after that, it should receive attention and this office must be contacted immediately. **Avoid physical activities during the first day,** as this can elevate your blood pressure and reinitiate bleeding.

- **Mouthcare:** You can start rinsing gently one day after the surgery. You may gently rinse your mouth with a solution of one teaspoon of salt dissolved in a 6oz. glass of warm (not hot) water. Avoid spitting as much as possible. Start using recommended mouthrinses very gently 2 days after the procedure, unless otherwise instructed by your periodontist. You may start brushing, flossing and using other dental aids *on other non-surgical areas* one day after the surgery. Do not brush, floss, or use other dental aids near the surgical site for 10 days. **Please avoid vigorous rinsing. Please do not disturb the surgical site by pulling your lip to look in your mouth.**

- **Discomfort and Swelling:** Following surgery, it is normal to experience some discomfort. You may be directed to gently apply ice packs to your face over the operated area for at least 3 – 4 hours following the surgery (with 15-min intervals). It may reduce the possibility of swelling and bruising in the following days. Maximum swelling may occur within 48 hours following surgery and may last for 5 – 7 days. At this time, you may use warm moist towel on your face. Pain may be more severe on the second or third day if there is swelling.

- **Medication:** Appropriate pain medication has been prescribed by your periodontist if necessary. Please use the prescribed medication as directed. **If you are prescribed antibiotics or narcotics, do not combine it with alcohol.** Do not drive or operate machinery while taking narcotic pain medications.

Please take the prescribed medication as instructed, unless adverse effects develop. Should skin rash, stomach problems, difficulty breathing, or other side effects occur, please stop taking your medication and contact the office immediately.

- **Numbness:** There may be numbness of your lip or cheek after the dental surgery. It will recover gradually. Please avoid biting on them. The doses and intervals of the prescribed pain medications are critical during the first couple of days. **It is important to start taking your pain medication before the numbness wears off.**

- **Diet:** Adequate food and fluid intake is important during the healing phase. You may start with cold drinks (cold water, iced tea, fruit juices without using straws) and soft diet for the first 2 – 3 days. A high-protein soft diet is desirable (e.g. yogurt, ice cream, cold soup, ground meat) during the healing phase. **Avoid hot and spicy foods during the healing phase.** Biting and chewing on the surgical side should be avoided as much as possible for 5 – 6 weeks. Advance the consistency of your diet gradually when you can chew comfortably.

- **Smoking:** Do not smoke. Refrain from smoking until the area is healed, and strongly consider quitting altogether.

- **Sutures:** Sutures will be removed by your periodontist in your post-op visit in 14 - 21 days. **Do not touch the surgical site or any loose sutures during the healing phase. The recommended stabilization tapes should be used for the first 2 – 3 weeks as instructed by your periodontist.**

Prolonged or excessive discomfort, swelling, bleeding or fever may seldom arise; Please contact the office immediately, so that further instructions for your care may be given.



## Origins Specialty Dentistry

5718 University Heights Blvd, Suite 201

San Antonio, TX 78249

Phone: (210) 460-0749 / Fax: (210) 560-2333

[info@originsspecialtydentistry.com](mailto:info@originsspecialtydentistry.com)

[www.OriginsSpecialtyDentistry.com](http://www.OriginsSpecialtyDentistry.com)